

hep-ex/9406001 2 Jun 94

LEP asymmetries and fits of the Standard Model

B. Pietrzyk

Laboratoire de Physique des Particules LAPP, IN2P3-CNRS,
F-74941 Annecy-le-Vieux, France

Abstract

The lepton and quark asymmetries measured at LEP are presented. The results of the Standard Model fits to the electroweak data presented at this conference are given. The top mass obtained from the fit to the LEP data is 172^{+13+18}_{-14-20} GeV; it is 177^{+11+18}_{-11-19} when also the collider, ν and A_{LR} data are included.

talk presented on the XXIXth Rencontres de Moriond
Méribel, March 12-19, 1994

It is a well known feature of the Standard Model that the interactions at the Z^0 resonance at LEP depend on the spin orientation of the incoming electrons and positrons as well as outgoing fermions. This feature creates the difference between the forward σ_F and backward σ_B cross-sections measured by the forward-backward asymmetries

$$A_{FB} = \frac{\sigma_F - \sigma_B}{\sigma_F + \sigma_B}$$

LEP experiments have measured the lepton asymmetries at different center of mass energies. The slope of asymmetries as a function of energy is inversely proportional to axial vector coupling constant squared g_A^2 or to the leptonic width Γ_l . The measured asymmetries are extrapolated to $\sqrt{s} = M_Z$, then the γ -, γZ -exchanges and the QED corrections are removed. We obtain in this way the $A^0 = \frac{3}{4}A_e A_f$ asymmetries, where

$$A_f = \frac{2g_V g_A}{g_V^2 + g_A^2} = \frac{2(1 - 4Q\sin^2\theta_w^{\text{eff}})}{1 + (1 - 4Q\sin^2\theta_w^{\text{eff}})^2}$$

The couplings and $\sin^2\theta_w^{\text{eff}}$ are "effective" since in their definition all possible loops in the Z self energy and the vertex corrections are included.

The lepton asymmetries measured by the LEP experiments are presented in Fig. 1. Contrary to the other experiments the L3 result was not updated with the 93 data. The LEP average is 0.0170 ± 0.0016 compared to 0.0161 ± 0.0019 during the summer conferences[1]. The lepton universality of the asymmetries is checked on Fig. 1. In the total error of 0.0016 the common systematical error of 0.0008 due to uncertainty of LEP energy scale is included. Also the noncorrelated systematical experimental error (typically 0.0006) is included. The theoretical error is not included. The difference between two LEP fitting programs ZFITTER 4.6 and MIZA(BHM) is 0.0008. Half of this difference comes from the treatment of initial state radiation ($O(\alpha^2)$ vs. YFS). The rest of the difference is under investigation.

The τ polarisation is measured by the difference between right-handed σ_R and left-handed σ_L cross-section for the τ production.

$$P_\tau(\cos\Theta) = \frac{\sigma_R - \sigma_L}{\sigma_R + \sigma_L} = -\frac{A_\tau(1 + \cos^2\Theta) + 2A_e\cos\Theta}{(1 + \cos^2\Theta) + 2A_eA_\tau\cos\Theta}$$

The mean τ polarisation measures A_τ while the forward-backward asymmetry of polarisation measures A_e . Typically the LEP experiments are measuring the angular distribution of the τ polarisation and obtain the A_e and A_τ from the fit of this distribution.

The results of the τ polarisation and the τ polarisation asymmetries are presented in Fig. 1. The mean values of A_e and A_τ from the τ polarisation give us A_l assuming lepton universality. The value of A_{LR} asymmetry from SLD [2], measuring also A_e , is given. There is a 1.9σ difference between A_e and A_τ measurements at LEP, 2.4σ between A_l and A_{LR} and 3.1σ between A_{LR} and A_e from τ polarisation at LEP. The LEP data up to 1992 have been analysed, the error is dominated by statistics, so improvement can be expected with already existing data.

The results on b and c asymmetries and for different techniques of measurements are presented on Fig. 2. Also for these measurements the statistical error dominates and not all the collected data are yet analysed. The b and c asymmetries presented in Fig.2 are asymmetries at $\sqrt{s}=91.3$ GeV and are corrected for QCD effects. We obtain the $A_{FB}^{0,b} = 0.096 \pm 0.004(stat) \pm 0.002(sys)$ and $A_{FB}^{0,c} = 0.070 \pm 0.008 \pm 0.007$ after applying the energy shift to

$\sqrt{s} = 91.19$ GeV, introducing the QED radiative corrections, removing the photon exchange and the γZ^0 -interference.

The lower part of Fig. 2 shows the values of $\sin^2 \theta_w^{\text{eff}}$ measured at LEP by different methods. The average value is 0.2322 ± 0.005 compared to 0.2321 ± 0.0006 reported during the 1993 summer conferences. It is interesting to note that the most precise measurement of $\sin^2 \theta_w^{\text{eff}}$ comes from forward-backward b asymmetries.

The result of the recent left-right asymmetry experiment at SLC is also combined with the LEP results. The $\chi^2/(d.o.f.)$ of the combined average is increasing from 6.3/5 to 12.8/6 when the SLC result is included. The difference between the SLC result and the combined LEP number is 2.5σ . The biggest difference, 3.1σ , is between the SLC left-right asymmetry and the τ forward-backward asymmetries. Both are measuring A_e and both errors are dominated by statistics.

The values of asymmetries as well as the other electroweak measurements presented at this conference are summarised in Table 1. They were used as input for the Standard Model fit. The results of the fit are presented in Table 2. A top mass of $172_{-14}^{+13+18}_{-20}$ is obtained from the fit to the LEP results, with an error improved since the 1993 summer conferences value of $166_{-19}^{+17+19}_{-22}$. The main improvement comes from the new Z width measurement. The central top mass value was obtained for the Higgs mass of 300 GeV and the second error describes the variation of the top mass with the Higgs mass between 60 and 1000 GeV. The α_s error of 0.005 becomes more and more interesting since the α_s measurement from R_l is one of the cleanest from the theoretical point of view. The $\chi^2/(d.o.f.)$ is 11.4/9 in comparison to 3.5/8 during summer conferences. A top mass of $177_{-11}^{+11+18}_{-19}$ is obtained when also the collider, ν -N and A_{LR} data are included in the fit. The A_{LR} SLC value is responsible for the top mass rise in the fit as well as for the increase of $\chi^2/(d.o.f.)$ to 19.1/12. The Standard Model fit parameters resulting from the fit including all data for the Higgs mass of 300 GeV are presented in Table 1 as well as the pulls. The forward-backward τ polarization asymmetries and R_b are pushing the top mass down with pulls of 1.8 and 2.0 respectively while the A_{LR} is driving it up with a pull of 2.7.

The $R_b = \Gamma_{b\bar{b}}/\Gamma_{\text{had}} = 0.2208 \pm 0.0024$ value is 2σ above the Standard Model prediction of 0.2158. The present value of R_b gives an upper limit on the top mass of 180 GeV independently of the Higgs mass. But if the difference of the R_b to the Standard Model persists it will have even more interesting consequences. It is interesting to follow this subject in the future. The R_b error of 0.0024 is obtained if the value of $\Gamma_{c\bar{c}}$ is left free in the full heavy flavour electroweak fit. If the value of R_c is fixed to 0.171 then R_b is 0.2207 ± 0.0022 . *

The comparison of the R_b , $\sin^2 \theta_w^{\text{eff}}$ and $R_\ell(R_{\text{had}})$ with the Standard Model predictions is presented in Figure 3. The deviation of R_b from the Standard Model can be seen on this plot. However R_ℓ is in agreement with the Standard Model for the $\alpha_s = 0.123 \pm 0.006$ used to produce the inclined band on this plot. For lower values of α_s the band moves up. It is seen from this plot that the present value of R_b favours the low top and low Higgs masses.

The results presented in this talk have been obtained by 4 LEP experiments and were averaged by the LEP Electroweak Working Group. I would like to thank D. Bardine, A. Blondel, J. Harton, S. de Jong, M. Martinez, M. Pepe-Altarelli, S. Rosier, D. Schaile, R. Tenchini and P. Wells for their help in preparation of this talk.

*In the Standard Model fit results presented here the value of R_b is slightly different from the value presented in my talk in Moriond due to some improvements in the averaging procedure. As a result the top mass is here slightly higher than in my Moriond presentation.

	measurement	Standard Model fit	pull
a) <u>LEP</u>			
line-shape and lepton asymmetries:			
M_Z [GeV]	91.1895 ± 0.0044	91.192	0.6
Γ_Z [GeV]	2.4969 ± 0.0038	2.4967	0.1
σ_h^0 [nb]	41.51 ± 0.12	41.44	0.6
R_ℓ	20.789 ± 0.040	20.781	0.2
$A_{\text{FB}}^{0,\ell}$	0.0170 ± 0.0016	0.0152	1.1
τ polarization:			
$\mathcal{A}_\tau$	0.150 ± 0.010	0.142	0.8
$\mathcal{A}_e$	0.120 ± 0.012	0.142	1.8
b and c quark results:			
$R_b = \Gamma_{b\bar{b}}/\Gamma_{\text{had}}$	0.2208 ± 0.0024	0.2158	2.0
$R_c = \Gamma_{c\bar{c}}/\Gamma_{\text{had}}$	0.170 ± 0.014	0.172	0.1
$A_{\text{FB}}^{0,b}$	0.0960 ± 0.0043	0.0997	0.8
$A_{\text{FB}}^{0,c}$	0.070 ± 0.011	0.071	0.1
$q\bar{q}$ charge asymmetry:			
$\sin^2\theta_{eff}^{\text{lept}}$ from $\langle Q_{\text{FB}} \rangle$	0.2320 ± 0.0016	0.2321	0.1
b) <u>p$\bar{p}$ and νN</u>			
M_W [GeV] (CDF, CDF prel., D0 prel., UA2; [3])	80.23 ± 0.18	80.31	0.4
$1 - M_W^2/M_Z^2(\nu N)$	0.2256 ± 0.0047	0.2246	0.2
c) <u>SLC</u>			
$\sin^2\theta_{eff}^{\text{lept}}$ from $\mathcal{A}_e$	0.2294 ± 0.0010	0.2321	2.7

Table 1: Summary of measurements included in the combined analysis of Standard Model parameters. Section a) summarizes LEP averages, section b) electroweak precision tests from hadron colliders and νN -scattering, section c) gives the result for $\sin^2\theta_{eff}^{\text{lept}}$ from the measurement of the left-right polarization asymmetry at SLC. The Standard Model fit result in column 3 and the pulls in column 4 are derived from the fit including all data (Table 2, column 4) for a fixed value of $M_H = 300$ GeV

References



- [1] The LEP Collaborations: ALEPH, DELPHI, L3 and OPAL, Phys. Lett. B276 (1992) 247; preprint CERN-PPE/93-157.
- [2] SLD Collaboration, K. Abe et al., SLAC-PUB-6456, submitted to Phys.Rev. Lett.
- [3] M. Demarteau et al., Combining W Mass Measurements, CDF/PHYS/CDF/PUBLIC/2552 and D0NOTE 2115.
- [4] UA2 Collaboration, J. Alitti et al., Phys. Lett. B276(1992) 354.

	LEP	LEP + Collider and ν data	LEP + Collider and ν data + A_{LR} from SLC
M_t (GeV)	$172^{+13}_{-14} \ ^{+18}_{-20}$	$170^{+12}_{-12} \ ^{+18}_{-19}$	$177^{+11}_{-11} \ ^{+18}_{-19}$
$\alpha_s(M_Z^2)$	$0.125 \pm 0.005 \pm 0.002$	$0.125 \pm 0.005 \pm 0.002$	$0.124 \pm 0.005 \pm 0.002$
$\chi^2/(d.o.f.)$	11.4/9	11.5/11	19.1/12
$\sin^2\theta_{eff}^{lept}$	$0.2323 \pm 0.0002 \ ^{+0.0001}_{-0.0002}$	$0.2324 \pm 0.0002 \ ^{+0.0001}_{-0.0002}$	$0.2320 \pm 0.0003 \ ^{+0.0001}_{-0.0002}$
$1 - M_W^2/M_Z^2$	$0.2251 \pm 0.0015 \ ^{+0.0003}_{-0.0003}$	$0.2253 \pm 0.0013 \ ^{+0.0003}_{-0.0002}$	$0.2243 \pm 0.0012 \ ^{+0.0003}_{-0.0002}$
M_W (GeV)	$80.28 \pm 0.08 \ ^{+0.01}_{-0.02}$	$80.26 \pm 0.07 \ ^{+0.01}_{-0.01}$	$80.31 \pm 0.06 \ ^{+0.01}_{-0.01}$

Table 2: Results of fits to LEP and other data for M_t and $\alpha_s(M_Z^2)$. No external constraint on $\alpha_s(M_Z^2)$ has been imposed. In the third column also the combined data from the $p\bar{p}$ experiments UA2 [4], CDF [5, 6] and D0 [6]: $M_W = 80.22 \pm 0.16$ GeV and from the neutrino experiments, CDHS [7], CHARM [8] and CCFR [9]: $1 - M_W^2/M_Z^2 = 0.2256 \pm 0.0047$ are included. The fourth column gives the result when also the SLD measurement of the left-right asymmetry at SLC [2], $\sin^2\theta_{eff}^{lept} = 0.2294 \pm 0.0010$, is added. The central values and the first errors quoted refer to $M_H = 300$ GeV. The second errors correspond to the variation of the central value when varying M_H in the interval $60 \leq M_H$ [GeV] ≤ 1000



- [5] CDF Collaboration, F. Abe et al., Phys. Rev. D43(1991) 2070.
- [6] CDF and D0 preliminary results, Y. Ducros, La Thuile, March 6-12, 1994.
- [7] CDHS Collaboration, H. Abramowicz et al., Phys. Rev. Lett. 57(1986) 298;
CDHS Collaboration, A. Blondel et al., Z.Phys. C45(1990) 361.
- [8] CHARM Collaboration, J.V. Allaby et al., Phys. Lett. B177(1986) 446;
CHARM Collaboration, J.V. Allaby et al., Z. Phys. C36(1987) 611.
- [9] CCFR Collaboration, NEVIS-R 1498, Submitted to Phys. Rev. Lett.
- [10] A. Blondel and C. Verzegnassi, Phys. Lett. B311(1993)346.

$A_{FB}^0(\text{lepton})$





A_{FB}^0

combined LEP data



τ Polarization $A_{\tau}^0 = -P_{\tau}$



τ Pol. Asymm. $A_e^0 = -4/3 \times P_{\tau}^{FB}$





Figure 1: Forward-backward lepton asymmetries, τ polarization and τ polarization asymmetries from LEP data.







Figure 2: Forward-backward b and c asymmetries and $\sin^2 \theta_W^{\text{eff}}$ measurements from LEP and SLC data.

SLC/LEP Preliminary APRIL 1994




Figure 3: The LEP/SLC measurements of $\sin^2 \theta_w^{eff}$ and R_b and the Standard Model predictions (updated figure from ref. [10]).